

Factors for delayed linkage to care following HIV diagnosis in the WHO European Region



Sara Croxford¹, Fiona Burns², Andrew Copas², Anastasia Pharris³, Valerie Delpech¹ for OptTEST by HIV in Europe

¹ Public Health England, HIV/STI Department, London, United Kingdom

² University College London, Infection & Population Health, London, United Kingdom

³ European Centre for Disease Prevention and Control, Stockholm, Sweden

BACKGROUND

Linking people who test HIV-positive to accessible and culturally appropriate care and support services is a crucial step in the HIV continuum of care.

Delayed linkage to HIV care is associated with delayed receipt of antiretroviral medications, faster disease progression and increased mortality.¹⁻⁴

Little comparable data are available on linkage to care in Europe.⁵

The objectives of this study were to utilise an existing surveillance dataset with a standardised definition to: i) describe linkage to HIV care following diagnosis in the World Health Organization (WHO) European Region and ii) to identify factors associated with delayed linkage.

METHODS

Analyses were carried out using data on new HIV diagnoses reported to the European Centre for Disease Prevention and Control (ECDC) and the WHO Regional Office for Europe in 2014. Data were included for countries that reported using the revised submission template (n=33/53).

Analyses were restricted to adults (aged ≥15 years) diagnosed from 2010-2014. Individuals were excluded if they had been previously diagnosed or in care, died within 3 months of diagnosis and/or had missing diagnosis/CD4 information.

All partial diagnosis, CD4 and death dates where only the month/quarter and year were reported, were defaulted to the middle of the month/quarter.

Logistic regression was used to determine factors for delayed linkage, adjusting for sex, age at diagnosis, diagnosis year, probable exposure, region of diagnosis, region of origin and first CD4 count after diagnosis.

Definitions

Linkage to care: patient seen for specialist HIV care after diagnosis (measured as the time between the HIV diagnosis date and first CD4 count date). The CD4 count date was used as a proxy for date of entry into HIV care.

Prompt linkage to care: patient seen for HIV care (CD4 count taken) in the 3 months (≤91 days) following diagnosis


Delayed linkage to care: patient seen for HIV care (CD4 count taken) more than 3 months (>91 days) after diagnosis

RESULTS

Of 125,665 adults diagnosed from 2010-2014, 3,794 were diagnosed previously, 769 were previously in HIV care and 810 died within three months of diagnosis.

A further 50,087 people had no CD4 data, 4,203 had incomplete CD4 or diagnosis dates (year only) and 5,863 had a CD4 count reported but no date (Figure 1).

Figure 1: Inclusion/exclusion criteria applied to calculate linkage to care



Among the 60,139 people included in these analyses, linkage to HIV care within 3 months of diagnosis was 95.5%. Prompt linkage by demographics and diagnosis characteristics can be seen in Figure 2.

In multivariable analysis, delayed linkage to care was associated with: being infected by injecting drug use or heterosexual transmission, being diagnosed in Central or Eastern Europe and having a first CD4 count >200 cells/mm³ (Figure 3).

Older age at diagnosis and being diagnosed after 2011 were associated with faster linkage to care. Sex and region of birth were not associated with linkage delays (Figure 3).

Figure 2: Prompt linkage to HIV care following diagnosis: Europe, 2010-2014





Figure 3: Multivariable analysis of factors associated with delayed linkage to care in Europe



CONCLUSIONS

Overall, linkage to care among adults diagnosed with HIV in Europe is prompt, with no difference in the timeliness of being linked by sex or region of birth.

However, given the high number of people with incomplete CD4 data, linkage estimates may be much lower than reported.

Our findings show improvements are needed in ensuring those diagnosed in Central and Eastern Europe and people infected through heterosexual contact and injecting drug use are promptly linked to care.

REFERENCES

- Ulett KB, Willig JH, Lin HY, et al. The therapeutic implications of timely linkage and early retention in HIV care. *AIDS Patient Care STDS* 2009 Jan;23(1):41-9.
- Mugavero MJ, Davila JA, Nevin CR, et al. From access to engagement: measuring retention in outpatient HIV clinical care. *AIDS Patient Care STDS* 2010 Oct;24(10):607-13.
- Gardner EM, McLees MP, Steiner JF, et al. The spectrum of engagement in HIV care and its relevance to test-and-treat strategies for prevention of HIV infection. *Clin Infect Dis* 2011 Mar 15;52(6):793-800.
- Croxford S, Kitching A, Desai S, Kall M, Edelstein M, Skingsley A, Burns F, Copas A, Brown AE, Sullivan AK, Delpech V. Mortality and causes of death in people diagnosed with HIV in the era of highly active antiretroviral therapy compared with the general population: an analysis of a national observational cohort. *Lancet Public Health* 2017; 2: e35-46
- Croxford S, Raben D, Burns, F, Delpech V for OptTEST by HIV in Europe. PE21/15 Linkage to care following HIV diagnosis in Europe: a review of the literature. *EAES* 2015.

OptTEST
PARTNERS

chip
Centre for
Health & Infectious
Disease Research

EA
TG European
AIDS Treatment
Group

REGION H Rigshospitalet

Aids
Fonds

MINISTERIO
DE ECONOMIA
Y COMPETITIVIDAD

i3i
Instituto de Salud
Carlos III

Public Health
England

Tervise Arengu Instituut
National Institute for Health Development

GNP+ GLOBAL NETWORK OF
PEOPLE LIVING WITH HIV

SSAT
ST STEPHEN'S AIDS TRUST

Inserm

MEDFASH
MEDICAL FOUNDATION FOR
HIV & SEXUAL HEALTH

We would also like to acknowledge the ECDC/WHO national focal points for their pivotal role in the surveillance of HIV in Europe



Co-funded by the 2nd Health
Programme of the
European Union



www.OptTEST.eu