



From Screening to Care - Coordinating Hepatologist

Overview

This case study from Hospital Universitari Vall d'Hebron in Barcelona in Spain evaluated an integrated viral hepatitis screening and linkage-to-care programme among emergency department patients requiring a blood test. The prevalence of both hepatitis B and hepatitis C was approximately three times higher than that reported in the general Spanish population.

Notably, most individuals diagnosed with viral hepatitis did not report any known risk factors. These findings suggest that risk-based screening alone may fail to identify a substantial proportion of infected individuals and support the value of routine screening approaches.

Approach

- HBV and HCV screening was integrated into the ED workflow and all individuals aged >18 requiring a blood test was offered hepatitis B and C testing.
- HCV antibodies and HBsAg were analysed using the same ED blood sample unless testing had been performed within the previous 3 months.
- Positive results were reviewed and validated by a hepatologist, who coordinated patient follow-up and referral to specialist care.
- Clinical records of patients with detectable HCV RNA or HBsAg were assessed to evaluate their clinical and social circumstances and guide management.
- Eligible patients were referred for specialist consultation, and those meeting treatment criteria were offered therapy at their first visit to minimise loss to follow-up.



Hepatitis C virus

0.7% HCV RNA-detectable (N = 128)

Men 46%, Median age 79 years

Bimodal distribution associated with age and sex



50-60 years old men
Psychiatric disorder



≥90 years old women
Prior blood transfusion



60% no risk factor identified



39% unaware of the infection



50% advanced fibrosis



Candidates for referral 53%

Linkage to care 65% (45/69)



Cost-effectiveness analysis:

1.06 QUALYs gained

ICUR 7,627 €/QUALYs

Participants 40-70 years: ICUR 767€/QUALYs



Hepatitis B virus

0.5% HBsAg positive (N = 91)



67% men

Median age: 62 years

23% Psychiatric disorder



92% no risk factor identified



41% unaware of the infection



35% no linked to care



Anti-HDV positive 2 cases



Candidates for referral 84%

Linkage to care 89% (42/46)



Cost-effectiveness analysis:

0.42 QUALYs gained

ICUR -147 €/QUALYs

Contry Case Study: Spain



EuroTEST


Working together for integrated
testing and earlier care

Addressing Hepatitis, HIV, STIs and TB

Lessons learned



In-hospital screening allows retrieval of medical records to help determine whether patients should be referred to a specialist.



Having a hepatologist act as the screening coordinator was considered a key factor in facilitating fast and simple linkage to care. This ensured that positive results didn't just remain within the usual testing cascade but were reviewed and followed up to the most appropriate care pathway



ED-based viral hepatitis screening was shown to be cost-effective for HCV and cost-saving for HBV, supporting its inclusion as part of broader hepatitis elimination strategies.



The programme has continued and been consolidated into routine clinical practice, with ongoing follow-up and data collection supporting continuous evaluation. This demonstrates a sustained institutional commitment and feasibility of maintaining ED-based viral hepatitis screening in a real-world setting.

Source

[Llaneras J, Ruiz-Cobo JC, Rando-Segura A, Barreira-Díaz A, Domínguez-Hernández R, Rodríguez-Frías F, Campins M, Colom J, Casado MA, Blanco-Grau A, Bañares J, Monforte A, Falcó-Roget A, Ruiz L, Meza B, Pumarola T, Riveiro-Barciela M, Esteban R, Buti M. Integrating viral hepatitis management into the emergency department: A further step towards viral hepatitis elimination. JHEP Rep. 2023 Oct 12;6\(1\):100932. doi: 10.1016/j.jhepr.2023.100932. PMID: 38074506; PMCID: PMC10698271.](#)

Interview with case study author

Please find below QR code for interview with study author, Jordi Llaneras, emergency physicians at Hospital Universitari Vall d'Hebron.

